

QUERY CONTROL FORM

RTIS USE ONLY

Application No. 09/182,753Prepared by Lois StoneTracking Number 5922421Examiner-GAU Bell - 1755Date 5/7/04Week Date 3/22/04No. of queries 1

IFW

JACKET

a. Serial No.	f. Foreign Priority	k. Print Claim(s)	p. PTO-1449
b. Applicant(s)	g. Disclaimer	l. Print Fig.	q. PTOL-85b
c. Continuing Data	h. Microfiche Appendix	m. Searched Column	r. Abstract
d. PCT	i. Title	n. PTO-270/328	s. Sheets/Figs
e. Domestic Priority	j. Claims Allowed	o. PTO-892	t. Other

SPECIFICATION

- a. Page Missing
- b. Text Continuity
- c. Holes through Data
- d. Other Missing Text
- e. Illegible Text
- f. Duplicate Text
- g. Brief Description
- h. Sequence Listing
- i. Appendix
- j. Amendments
- k. Other

CLAIMS

- a. Claim(s) Missing
- b. Improper Dependency
- c. Duplicate Numbers
- d. Incorrect Numbering
- e. Index Disagrees
- f. Punctuation
- g. Amendments
- h. Bracketing
- i. Missing Text
- j. Duplicate Text
- k. Other

MESSAGE


There are two original claims numbered 32.
One is attached to the end of original claim 31.
Copy provided for reference.

Thanks you,

initials LS

RESPONSE

initials




wherein Cp_a is a substituted cyclopentadienyl ring, Cp_b is an unsubstituted or substituted cyclopentadienyl ring; each R is the same or different and is a hydrocarbyl radical having 1-20 carbon atoms; each R'_m is the same or different and is a hydrocarbyl radical having 1-20 carbon atoms; R'' is a structural bridge between the cyclopentadienyl rings imparting stereorigidity to the metallocene and is selected from the group consisting of an alkylene radical having 1-4 carbon atoms, a silicon hydrocarbyl radical, a germanium hydrocarbyl radical, a phosphorus hydrocarbyl radical, a nitrogen hydrocarbyl radical, a boron hydrocarbyl radical, and an aluminum hydrocarbyl radical; Me is a group 4b, 5b, or 6b metal from the Periodic Table of Elements; each Q is a hydrocarbyl radical having 1-20 carbon atoms or is a halogen; $0 \leq p \leq 3$; $0 \leq m \leq 4$; $1 \leq n \leq 4$; and

wherein R'_m is selected such that $(Cp_bR'_m)$ is a sterically different ring than (Cp_aR_n) . ³² The

process of claim 31 wherein the weight ratio of said alumoxane to said silica is within the range of about 0.7-1.0. ←

³⁰ ~~32~~. (original) The process of claim ²⁸~~29~~ wherein the weight ratio of said alumoxane to said silica is within the range of about 0.7-1.0. ←

³¹ ~~33~~. (currently amended) The ~~method~~ process of claim ²⁶~~27~~ wherein said stereospecific metallocene is an isospecific stereorigid metallocene characterized by the formula:



wherein each $(C_5(R')_4)$ is a substituted cyclopentadienyl ring; each R' is the same or different and is a hydrogen or hydrocarbyl radical having 1-20 carbon atoms; R'' is a structural bridge between the two $(C_5(R')_4)$ rings imparting stereorigidity to said metallocene with the two $(C_5(R')_4)$ rings being in